




KINETEX[®]
Core-Shell Technology

NEW Kinetex **F5**

HPLC/UHPLC Core-Shell Columns

Whoa! I can
even separate
structural
isomers!



- Reduce Method Development Time by Days
- Greater Reproducibility than other **PFPs**
- 5 Glorious Interaction Mechanisms
- 5 Valuable LC Separation Modes

F5

TM


phenomenex[®]
...breaking with traditionSM



How I Work

With the astonishing combination of core-shell performance and 5 interaction mechanisms, Kinetex® F5 columns will effortlessly drive your orthogonal HPLC/UHPLC development!

- A Hydrophobic**
Carbon skeleton of linker and ring encourage neutral/hydrophobic retention
- B Aromatic**
In non-acetonitrile mobile phases, π - π electrons of the carbon ring interact with analyte π - π electrons and result in positive retention increase



- C Electrostatic**
High electronegativity of the fluorine groups create dipole moments, aiding in polar compound retention. Induced dipole moments can also aid neutral compound retention.
- D Steric/Planar**
Shape selectivity allows for isomeric separations that are otherwise impossible on traditional alkyl phases.
- E Hydrogen Bonding**
Extremely effective retention mechanism caused as polar functional groups of analyte interact with the electron greedy fluorine.

Why I'm Better

While older pentafluorophenyl phases (PFP, PFPP, F5, etc.) are based on existing bonding techniques and technologies that promote irreproducibility, the Kinetex F5 was meticulously designed by Phenomenex R&D and its customers, to provide consistently accurate and high performance results.


Learn how the new Kinetex F5 will get you the results you deserve time and time again!



- Core-Shell Advantage **p. 4**
- Method Development Versatility **p. 5**
- Unmatched Reproducibility **pp. 6-7**
- 100% Aqueous Stability **p. 8**
- Isomeric Separations **p. 9**
- Vitamin D3 Epimers **p. 10**
- Multi Component Screening **p. 11**
- Methanol vs. Acetonitrile **p. 12**
- Trace Impurity Detection **p. 13**
- UHPLC Consistency
and Performance **pp. 14-15**
- Kinetex Column Selection **pp. 16-17**
- Ordering Information **pp. 18-19**



If you are not completely satisfied with Kinetex core-shell columns, send in your comparative data to a similar product with the Kinetex column within 45 days for a FULL REFUND.




Why Wait?


You Can Get Improvements Today!

The band broadening (wide peaks) and lengthy retention times of traditional fully porous products can be limiting your results. Now you can use the incredible efficiency levels of Kinetex® Core-Shell Technology to achieve shorter run times, higher levels of sensitivity, and overall better HPLC/UHPLC results.

Core-Shell vs. Fully Porous Efficiency Levels (plates/m)



Core-Shell Performance Gains



■ Kinetex Core-Shell
■ Fully Porous


Get faster,
better results
with me!



Method Development

Versatility

Combine core-shell performance, multiple retention mechanisms and the Kinetex F5 column's ability to be run in a variety of separation modes and you now have an impeccable method development tool at your disposal.






Dependability




Batch-to-Batch, Column-to-Column

Conventional fully porous and core-shell PFP/F5 columns fail to reach the level of repeatability that you deserve. Inconsistencies in their base silica have led to data inaccuracies that waste your time and money. Kinetex® F5 columns were specifically designed to avoid these past problems and provide you with the highest degree of reproducibility.

Kinetex 2.6µm F5



Supelco® Ascentis® Express 2.7µm F5



Conditions for all columns:

Column: Kinetex 2.6µm F5
 Discovery 3µm HS F5
 Ascentis Express 2.7µm F5
 Hypersil GOLD 3µm PFP
 Accucore 2.6µm PFP

Dimensions: 50 x 4.6 mm

Mobile Phase: A: 0.1 % Formic acid in Water
 B: 0.1 % Formic acid in Acetonitrile

Gradient: 5-95 % B over 5 minutes

Flow Rate: 1.85 mL/min

Temperature: Ambient


Detection: UV @ 254 nm

Sample:
 1. Uracil
 2. Pindolol
 3. Chlorpheniramine
 4. Nortriptyline
 5. 3-Methyl-4-Nitrobenzoic acid
 6. 5-Methyl Salicyl Aldehyde
 7. Hexaphenone


What happened to peaks 3 and 4?

Accucore is a trademark and Hypersil GOLD is registered trademark of Thermo Hypersil-Keystone. Ascentis, Discovery and Supelco are registered trademarks of Sigma-Aldrich Co. LLC. Phenomenex is not affiliated with any of the above companies. Comparative separations may not be representative of all applications.

Supelco® Discovery® 3µm HS F5




Thermo Hypersil GOLD® 3µm PFP



Adsorption of Peaks 3 and 4!

Retention Time Shifting

Thermo Accucore™ 2.6µm PFP



Conditions for all columns:

Column: Kinetex 2.6µm F5
Discovery 3µm HS F5
Ascentis Express 2.7µm F5
Hypersil GOLD 3µm PFP
Accucore 2.6µm PFP

Dimensions: 50 x 4.6 mm

Mobile Phase: A: 0.1% Formic acid in Water
B: 0.1% Formic acid in Acetonitrile

Gradient: 5-95% B over 5 minutes

Flow Rate: 1.85 mL/min


Temperature: Ambient

Detection: UV @ 254 nm

Sample: 1. Uracil
2. Pindolol
3. Chlorpheniramine
4. Nortriptyline
5. 3-Methyl-4-Nitrobenzoic acid
6. 5-Methyl Salicyl Aldehyde
7. Hexaphenone

Are you okay with peak adsorptions and retention time shifts?




Accucore is a trademark and Hypersil GOLD is registered trademark of Thermo Hypersil-Keystone. Ascentis, Discovery and Supelco are registered trademarks of Sigma-Aldrich Co. LLC. Phenomenex is not affiliated with any of the above companies. Comparative separations may not be representative of all applications.



Extra Polar Retention and 100% Aqueous Stability



Unlike traditional alkyl stationary phases, the polar functional group of the Kinetex® F5 makes it stable in 100% aqueous mobile phase conditions. This can be especially beneficial for methods that require the retention and resolution of problematic polar compounds that typically have poor retention under reversed phase conditions.

Organic Acids



Each successive day, column was equilibrated, run and stored in the 100% aqueous conditions mentioned above.

Column: Kinetex 2.6µm F5
Dimensions: 100 x 4.6mm
Part No.: 00D-4723-E0
Mobile Phase: 20 mM Sodium phosphate pH 2.5
Flow Rate: 1.5 mL/min
Temperature: Ambient
Detection: UV @ 210 nm
Sample: 1. Tartaric Acid
2. Malic Acid
3. Maleic Acid
4. Fumaric Acid
5. Succinic Acid




INCREDIBLE!
Even the isomers
Maleic acid and
Fumaric acid
are separated!



Novel Selectivity For Isomeric Separations

While a C18 can differentiate between the small addition of a single methyl group between two similar compounds, it cannot separate compounds with only structural differences, like positional isomers. This is where the electrostatic and planar interactions of the Kinetex F5 give focused attention to the resolution you need.

Methoxybenzene Isomers



Conditions for all columns:

Column: Kinetex 2.6µm F5
Kinetex 2.6µm C18
Kinetex 2.6µm Biphenyl

Dimensions: 150 x 4.6 mm

Mobile Phase: A: 0.1% TFA in Water
B: Acetonitrile



Isocratic: A/B (65:35)

Flow Rate: 1 mL/min

Temperature: Ambient

Detection: UV @ 254 nm

Sample: 1. 1,2,3-Trimethoxybenzene
2. 1,2-Dimethoxybenzene
3. 1,2,4-Trimethoxybenzene
4. 1,4-Dimethoxybenzene
5. Methoxybenzene
6. 1,3-Dimethoxybenzene




Clinical Research

Vitamin D3 Epimers

Even tandem mass spectrometry (LC/MS/MS) analysis has need to utilize the wonderful cross functionality of Kinetex® F5. With the same fragment ions coming from the Vitamin D3 epimers, reproducible chromatographic separation is a must. The unique combination of polar/non-polar selectivity of Kinetex F5 easily performs this necessary separation in a highly sensitive and short analysis window.


Kinetex 2.6 µm F5




Conditions for all columns:

- Column:** Kinetex 2.6 µm F5
HALO 2.7 µm PFP
XSelect HSS 2.5 µm PFP
- Dimensions:** 100 x 4.6 mm
- Mobile Phase:** A: 0.1% Formic acid in Water
B: 0.1% Formic acid in Methanol
- Isocratic:** A/B (15:85)
- Flow Rate:** 0.75 mL/min
- Temperature:** Ambient
- Detection:** MS/MS (AB SCIEX API 4000™)
- Sample:** 1. 25-OH Vitamin D3
2. 25-OH Vitamin D2
3. 3-epi-25-OH Vitamin D3

Advanced Material Technology HALO® 2.7 µm PFP



Waters® XSelect® HSS 2.5 µm PFP



Vitamin D3 epimers are not fully separated


HALO is a registered trademark of Advanced Materials Technology, Inc. Waters and XSelect are registered trademarks of Waters Corporation. Phenomenex is not affiliated with any of the above companies. Comparative separations may not be representative of all applications.

Food Testing

Multi Component Analysis


The versatility of Kinetex F5 core-shell columns matches marvelously with ingredient quantification and contaminant identification. Acidic food additives may tail and lack retention on a C18 column, but the Kinetex F5 offers superb polar retention and performance. Meanwhile, large multi component contaminant screens and their mixtures of acids, bases, neutrals and structurally similar compounds, can easily utilize the combination of polar, non-polar and geometric interaction mechanisms of the Kinetex F5 to get excellent separation and sensitivity.

Food Additives



Column: Kinetex 2.6µm F5
Dimensions: 150 x 2.1 mm
Part No.: 00F-4723-AN
Mobile Phase: A: 0.1% Phosphoric acid in Water
 B: Acetonitrile
Gradient: 5-35% A in 6 min. Hold for 4 min.
Flow Rate: 0.6 mL/min
Temperature: 30 °C
Detection: UV @ 240 nm
Sample: 1. Saccharin
 2. p-Hydroxybenzoic acid
 3. Sorbic acid
 4. Dehydroacetic acid
 5. Methyl paraben

Antibiotics Screen



Column: Kinetex 2.6µm F5
Dimensions: 50 x 2.1 mm
Part No.: 00B-4723-AN
Mobile Phase: A: 0.1% Formic acid in Water
 B: 0.1% Formic acid in Methanol

Gradient:	Time (min)	% B
	0	0
	0.08	0
	1.08	100
	8.08	100
	11.08	0
	11.58	0

Flow Rate: 0.5 mL/min
Temperature: 30 °C
Detection: MS/MS (AB SCIEX API 4000™)
Sample: 1. Amoxicillin
 2. Cefalexin
 3. Cefazolin
 4. Cefoperazone
 5. Cefapirin
 6. Cloxacillin
 7. Dicloxacillin
 8. Ciprofloxacin
 9. Difloxacin
 10. Marbofloxacin
 11. Sulfadiazine
 12. Sulfamerazine
 13. Sulfamethazine
 14. Sulfamethoxazole
 15. Sulfapyridine
 16. Sulfaquinoxaline
 17. Sulfathiazole
 18. Neospiramycin
 19. Doxycycline
 20. Tiamulin
 21. Valnemulin
 22. Rifaximin
 23. Lincomycin
 24. Nafcillin

Find more Kinetex F5 applications at www.phenomenex.com/KinetexF5Apps





Methanol vs. Acetonitrile

A Phenyl Story

While mobile phase modifiers can help adjust retention, with the use of the Kinetex® F5, the major mobile phase constituent can also be used to manipulate elution order and retention properties. Acetonitrile can be used to disrupt pi-pi interactions between compounds and phenyl phases, while switching to the weaker solvent Methanol will encourage aromatic interactions.

Methoxybenzene Isomers



Conditions for all columns:

Column: Kinetex 2.6 μ m F5

Dimensions: 150 x 4.6 mm

Part No.: 00F-4723-E0

Mobile Phase: as noted



Isocratic: A/B (65:35)

Flow Rate: 1 mL/min

Temperature: Ambient

Detection: UV @ 254 nm

Sample: 1. 1,2,3-Trimethoxybenzene
2. 1,2-Dimethoxybenzene
3. 1,2,4-Trimethoxybenzene




Impact selectivity and retention with solvent adjustment!


Trace Pharmaceutical Impurity Detection

Trace impurities of active pharmaceutical ingredients are incredibly important to identify and quantify. With the rapid performance value of core-shell technology combined with the versatility of a pentafluorophenyl, the Kinetex F5 is the precise alternative to other reversed phase columns that you need. Easily utilize the Kinetex F5 to get greater sensitivity, better resolution and all in shorter analysis times.

Kinetex 1.7 μ m F5



Waters® ACQUITY® CSH™ 1.7 μ m Fluoro-Phenyl



Conditions for all columns:

Column: Kinetex 1.7 μ m F5
ACQUITY CSH 1.7 μ m Fluoro-Phenyl

Dimensions: 50 x 2.1 mm

Mobile Phase: A: 20mM Potassium phosphate pH 2.3
B: Methanol

Gradient:	Time (min)	% B
	0	5
	10	95
	10.01	5

Flow Rate: 0.3 mL/min

Temperature: Ambient


Detection: UV @ 254 nm

Sample:
1. Impurity 1
2. Impurity 2
3. Proprietary Active Pharmaceutical Ingredient



Incredible UHPLC Reproducibility

Highly sensitive UHPLC separations are dependent upon consistent quantitation and consistent results. With the reliability of the Kinetex® F5, you no longer have to settle for the inconsistent results of PFP products that currently exist on the market. Our highest standards of quality will ensure that you are fully satisfied with each and every Kinetex F5 column!

Trusted Consistency



Column: Kinetex 1.7 μ m F5
Dimensions: 50 x 4.6 mm
Mobile Phase: A: 0.1 % Formic in Water
B: 0.1 % Formic in Acetonitrile
Gradient: 5-95 % B over 5 min.
Flow Rate: 1.85 mL/min
Temperature: Ambient
Detection: UV @ 254 nm
Sample:
1. Uracil
2. Pindolol
3. Chlorpheniramine
4. Nortriptyline
5. 3-Methyl-4-Nitrobenzoic acid
6. 5-Methyl Salicyl Aldehyde
7. Hexaphenone




A Better PFP

Selectivity for UHPLC

Kinetex 1.7 μm core-shell technology produces increased efficiencies over traditional sub-2 μm columns on the market, yielding remarkable chromatographic resolution, higher peak capacities, and greater sensitivity. Add these performance gains alongside the novel selectivity and excellent reproducibility of the Kinetex F5 and you now have an incredible UHPLC solution at your fingertips.

Pharmaceutical Mix



Conditions for all columns:

Column: Kinetex 1.7 μm F5
ACQUITY CSH 1.7 μm Fluoro-Phenyl
ACQUITY HSS 1.8 μm PFP

Dimensions: 50 x 2.1 mm

Mobile Phase: A: 10 mM Ammonium acetate pH 3.2
B: Acetonitrile



Gradient	Time (min)	% B
	90	10
	90	10
	5	95
	5	95
	90	10
	90	10

Flow Rate: 0.4 mL/min

Temperature: Ambient

Detection: UV @ 220 nm

Sample:
1. Pindolol
2. Propranolol
3. Indoprofen
4. Naproxen
5. Warfarin
6. Terfenadine



Terrellasaurus – first discovered in Torrance, CA.

Waters and ACQUITY are registered trademarks, and CSH is a trademark of Waters Corporation. Phenomenex is not affiliated with Waters Corporation. Comparative separations may not be representative of all applications.

Get Method Development Flexibility For Your Small Molecule Analysis

Recommended Selectivities If You're Working With:

Acids

- Kinetex® C18
- Kinetex F5
- Kinetex Phenyl-Hexyl

Bases

- Kinetex EVO C18
- Kinetex XB-C18
- Kinetex Biphenyl

Neutrals

- Kinetex C18
- Kinetex C8
- Kinetex Biphenyl

Aromatics

- Kinetex Biphenyl
- Kinetex Phenyl-Hexyl
- Kinetex F5

Acids, Bases, and Neutrals

- Kinetex C18
- Kinetex Biphenyl
- Kinetex EVO C18
- Kinetex F5

Highly Polar Compounds

- Kinetex EVO C18
- Kinetex F5
- Kinetex Biphenyl
- Kinetex HILIC

High pH

- Kinetex EVO C18

Isomers

- Kinetex F5

Upgrading Your Fully Porous Methods:

Fully Porous 3 μm or 5 μm

- **Kinetex 5 μm** – Drop-in for easy performance improvements with no backpressure increase
- **Kinetex 2.6 μm** – Dramatically improve results with efficiency/peak capacity gains

Fully Porous sub-2 μm

- **Kinetex 2.6 μm** – Get similar efficiencies at lower backpressure allowing for greater productivity gains
- **Kinetex 1.7 μm** – Up to 20% greater efficiencies resulting in drop-in improvements
- **Kinetex 1.3 μm** – Incredible efficiency gains on high end UHPLC systems

Fully Porous Preparative LC

- **Kinetex 5 μm** – Drop-in for easy performance improvement with no backpressure increase

Choosing The Best Core-Shell Platform For You is Easy!

For Small Molecules

		5 μ m	3.6 μ m	2.6 μ m	1.7 μ m	1.3 μ m
UHPLC						
HPLC						
PREP LC						

Phase	Best Use	pH Stability	Available Particle Size(s)			
F5	Highly reproducible pentafluorophenyl propyl phase that offers a unique combination of polar, hydrophobic, aromatic, and shape selectivity	1.5 - 8.5*		2.6 μ m	1.7 μ m	
EVO C18	Robust reversed phase methods even in alkaline conditions with improved peak shape for polar basic compounds	1 - 12	5 μ m			
C18	All purpose phase that offers the hydrophobic retention and methylene selectivity chromatographers expect from a C18 column	1.5 - 8.5*	5 μ m	2.6 μ m	1.7 μ m	1.3 μ m
XB-C18	C18 phase with protective butyl side chains for improved peak shape for basic compounds under neutral and acidic conditions	1.5 - 8.5*	5 μ m	2.6 μ m	1.7 μ m	
C8	USP L7 phase that provides less hydrophobic and methylene selectivity than a C18	1.5 - 8.5*	5 μ m	2.6 μ m	1.7 μ m	
Biphenyl	100% aqueous stable and allows for excellent reversed phase retention and enhanced polar and aromatic selectivity	1.5 - 8.5*	5 μ m	2.6 μ m	1.7 μ m	
Phenyl-Hexyl	Reversed phase chemistry that allows for greater retention and separation of aromatic hydrocarbons	1.5 - 8.5*	5 μ m	2.6 μ m	1.7 μ m	
HILIC	Unbonded silica phase for HILIC conditions to provide selectivity for polar compounds	2.0 - 7.5	5 μ m	2.6 μ m	1.7 μ m	

Phenomenex Application Specific Core-Shell Products

Material	Phase	Best Use	pH Stability	Available Particle Size(s)			
For Peptides (\leq 10,000 Da)							
Aeris™ PEPTIDE	XB-C18	Excellent hydrophobicity and methylene selectivity for peptide and peptide mapping separations	1.5 - 9.0	5 μ m	3.6 μ m	2.6 μ m	1.7 μ m
For Proteins (> 10,000 Da)							
Aeris WIDEPORE	XB-C18	Maximum hydrophobicity and high temp stability for hydrophilic and PEGylated proteins	1.5 - 9.0		3.6 μ m		
	XB-C8	Medium hydrophobicity and high temp stability for moderately hydrophobic proteins and glycosylated proteins	1.5 - 9.0		3.6 μ m		
	C4	Lowest hydrophobicity for very large or very hydrophobic proteins	1.5 - 9.0		3.6 μ m		
For Synthetic Oligonucleotides (DNA/RNA)							
Clarity® Oligo-MS™	C18	Rapid, high efficiency reversed phase LC/MS analysis for QC and characterization	1.5 - 8.5*			2.6 μ m	1.7 μ m

*pH stability under gradient conditions. pH stability is 1.5-10 under isocratic conditions.

Ordering Information

5 µm Minibore Columns (mm)					SecurityGuard™ ULTRA Cartridges‡
Phases	30 x 2.1	50 x 2.1	100 x 2.1	150 x 2.1	3/pk
EVO C18	00A-4633-AN	00B-4633-AN	00D-4633-AN	00F-4633-AN	AJ0-9298
Biphenyl	00A-4627-AN	00B-4627-AN	00D-4627-AN	—	AJ0-9209
XB-C18	00A-4605-AN	00B-4605-AN	00D-4605-AN	—	AJ0-8782
C18	00A-4601-AN	00B-4601-AN	00D-4601-AN	00F-4601-AN	AJ0-8782
C8	—	00B-4608-AN	00D-4608-AN	—	AJ0-8784
Phenyl-Hexyl	—	00B-4603-AN	00D-4603-AN	—	AJ0-8788

for 2.1 mm ID

5 µm MidBore™ Columns (mm)				SecurityGuard ULTRA Cartridges‡
Phases	50 x 3.0	100 x 3.0	150 x 3.0	3/pk
EVO C18	00B-4633-Y0	00D-4633-Y0	00F-4633-Y0	AJ0-9297
Biphenyl	00B-4627-Y0	00D-4627-Y0	00F-4627-Y0	AJ0-9208
XB-C18	00B-4605-Y0	00D-4605-Y0	00F-4605-Y0	AJ0-8775
C18	00B-4601-Y0	00D-4601-Y0	00F-4601-Y0	AJ0-8775
C8	00B-4608-Y0	00D-4608-Y0	—	AJ0-8777
Phenyl-Hexyl	00B-4603-Y0	00D-4603-Y0	—	AJ0-8781

for 3.0 mm ID

5 µm Analytical Columns (mm)					SecurityGuard ULTRA Cartridges‡
Phases	50 x 4.6	100 x 4.6	150 x 4.6	250 x 4.6	3/pk
EVO C18	00B-4633-E0	00D-4633-E0	00F-4633-E0	00G-4633-E0	AJ0-9296
Biphenyl	00B-4627-E0	00D-4627-E0	00F-4627-E0	00G-4627-E0	AJ0-9207
XB-C18	00B-4605-E0	00D-4605-E0	00F-4605-E0	00G-4605-E0	AJ0-8768
C18	00B-4601-E0	00D-4601-E0	00F-4601-E0	00G-4601-E0	AJ0-8768
C8	00B-4608-E0	00D-4608-E0	00F-4608-E0	00G-4608-E0	AJ0-8770
Phenyl-Hexyl	00B-4603-E0	00D-4603-E0	00F-4603-E0	00G-4603-E0	AJ0-8774

for 4.6 mm ID

5 µm Semi-Preparative Columns (mm)			SecurityGuard SemiPrep Cartridges***
Phases	150 x 10	250 x 10	10 x 10 /3pk
C18	00F-4601-N0	00G-4601-N0	AJ0-9278
Biphenyl	00F-4627-N0	00G-4627-N0	AJ0-9280

for 10 mm ID

5 µm Axia™ Packed Preparative Columns (mm)					SecurityGuard PREP Cartridges*
Phases	50 x 21.2	100 x 21.2	150 x 21.2	250 x 21.2	15 x 21.2
EVO C18	00B-4633-P0-AX	00D-4633-P0-AX	00F-4633-P0-AX	00G-4633-P0-AX	AJ0-9304 /ea
Biphenyl	00B-4627-P0-AX	00D-4627-P0-AX	00F-4627-P0-AX	00G-4627-P0-AX	AJ0-9272 /ea
XB-C18	00B-4605-P0-AX	00D-4605-P0-AX	00F-4605-P0-AX	00G-4605-P0-AX	AJ0-9145
C18	00B-4601-P0-AX	00D-4601-P0-AX	00F-4601-P0-AX	00G-4601-P0-AX	AJ0-9145
C8	00B-4608-P0-AX	00D-4608-P0-AX	00F-4608-P0-AX	00G-4608-P0-AX	AJ0-9205
Phenyl-Hexyl	00B-4603-P0-AX	00D-4603-P0-AX	00F-4603-P0-AX	00G-4603-P0-AX	AJ0-9147
HILIC	—	00D-4606-P0-AX	00F-4606-P0-AX	00G-4606-P0-AX	AJ0-9277

for 21.2 mm ID

5 µm Axia Packed Preparative Columns (mm)					SecurityGuard PREP Cartridges**
Phases	50 x 30	100 x 30	150 x 30	250 x 30	15 x 30
EVO C18	—	00D-4633-U0-AX	00F-4633-U0-AX	00G-4633-U0-AX	AJ0-9305 /ea
Biphenyl	—	—	00F-4627-U0-AX	—	AJ0-9273
XB-C18	00B-4605-U0-AX	00D-4605-U0-AX	00F-4605-U0-AX	00G-4605-U0-AX	AJ0-9204
C18	00B-4601-U0-AX	00D-4601-U0-AX	00F-4601-U0-AX	00G-4601-U0-AX	AJ0-9204
C8	00B-4608-U0-AX	00D-4608-U0-AX	00F-4608-U0-AX	00G-4608-U0-AX	AJ0-9217
Phenyl-Hexyl	00B-4603-U0-AX	00D-4603-U0-AX	00F-4603-U0-AX	00G-4603-U0-AX	AJ0-9216

for 30 mm ID



Remove interferences such as proteins, phospholipids, salts, and more in a quick and simple procedure
www.phenomenex.com/Novum



Convenient and inexpensive way to remove particulates from HPLC samples
www.phenomenex.com/Phenex



Protect your Kinetex LC columns from contaminants and extend their lifetime.
www.phenomenex.com/guardit



If you are not completely satisfied with Kinetex core-shell columns, send in your comparative data to a similar product with the Kinetex column within 45 days for a FULL REFUND.

† SecurityGuard ULTRA Cartridges require holder, Part No.: AJ0-9000

* PREP SecurityGuard Cartridges require holder, Part No.: AJ0-8223

** PREP SecurityGuard Cartridges require holder, Part No.: AJ0-8277

*** SemiPrep SecurityGuard Cartridges require holder, Part No.: AJ0-9281

Ordering Information

2.6 µm Minibore Columns (mm)						SecurityGuard™ ULTRA Cartridges [‡]
Phases	30 x 2.1	50 x 2.1	75 x 2.1	100 x 2.1	150 x 2.1	3/pk
F5	—	00B-4723-AN	—	00D-4723-AN	00F-4723-AN	AJO-9322
Biphenyl	00A-4622-AN	00B-4622-AN	—	00D-4622-AN	00F-4622-AN	AJO-9209
XB-C18	00A-4496-AN	00B-4496-AN	00C-4496-AN	00D-4496-AN	00F-4496-AN	AJO-8782
C18	00A-4462-AN	00B-4462-AN	00C-4462-AN	00D-4462-AN	00F-4462-AN	AJO-8782
C8	00A-4497-AN	00B-4497-AN	00C-4497-AN	00D-4497-AN	00F-4497-AN	AJO-8784
HILIC	00A-4461-AN	00B-4461-AN	00C-4461-AN	00D-4461-AN	00F-4461-AN	AJO-8786
Phenyl-Hexyl	00A-4495-AN	00B-4495-AN	00C-4495-AN	00D-4495-AN	00F-4495-AN	AJO-8788

for 2.1 mm ID

2.6 µm MidBore™ Columns (mm)						SecurityGuard™ ULTRA Cartridges [‡]
Phases	30 x 3.0	50 x 3.0	75 x 3.0	100 x 3.0	150 x 3.0	3/pk
F5	—	00B-4723-YO	—	00D-4723-YO	00F-4723-YO	AJO-9321
Biphenyl	—	00B-4622-YO	—	00D-4622-YO	00F-4622-YO	AJO-9208
XB-C18	00A-4496-YO	00B-4496-YO	00C-4496-YO	00D-4496-YO	00F-4496-YO	AJO-8775
C18	00A-4462-YO	00B-4462-YO	00C-4462-YO	00D-4462-YO	00F-4462-YO	AJO-8775
C8	00A-4497-YO	00B-4497-YO	00C-4497-YO	00D-4497-YO	00F-4497-YO	AJO-8777
HILIC	00A-4461-YO	—	—	—	00F-4461-YO	AJO-8779
Phenyl-Hexyl	—	00B-4495-YO	—	00D-4495-YO	00F-4495-YO	AJO-8781

for 3.0 mm ID

2.6 µm Analytical Columns (mm)						SecurityGuard™ ULTRA Cartridges [‡]
Phases	30 x 4.6	50 x 4.6	75 x 4.6	100 x 4.6	150 x 4.6	3/pk
F5	—	00B-4723-E0	—	00D-4723-E0	00F-4723-E0	AJO-9320
Biphenyl	—	00B-4622-E0	—	00D-4622-E0	00F-4622-E0	AJO-9207
XB-C18	—	00B-4496-E0	00C-4496-E0	00D-4496-E0	00F-4496-E0	AJO-8768
C18	00A-4462-E0	00B-4462-E0	00C-4462-E0	00D-4462-E0	00F-4462-E0	AJO-8768
C8	—	00B-4497-E0	00C-4497-E0	00D-4497-E0	00F-4497-E0	AJO-8770
HILIC	—	00B-4461-E0	00C-4461-E0	00D-4461-E0	00F-4461-E0	AJO-8772
Phenyl-Hexyl	—	00B-4495-E0	00C-4495-E0	00D-4495-E0	00F-4495-E0	AJO-8774

for 4.6 mm ID

1.7 µm Minibore Columns (mm)					SecurityGuard™ ULTRA Cartridges [‡]
Phases	30 x 2.1	50 x 2.1	100 x 2.1	150 x 2.1	3/pk
F5	—	00B-4722-AN	00D-4722-AN	00F-4722-AN	AJO-9322
Biphenyl	—	00B-4628-AN	00D-4628-AN	00F-4628-AN	AJO-9209
XB-C18	00A-4498-AN	00B-4498-AN	00D-4498-AN	00F-4498-AN	AJO-8782
C18	00A-4475-AN	00B-4475-AN	00D-4475-AN	00F-4475-AN	AJO-8782
C8	00A-4499-AN	00B-4499-AN	00D-4499-AN	00F-4499-AN	AJO-8784
HILIC	00A-4474-AN	00B-4474-AN	00D-4474-AN	—	AJO-8786
Phenyl-Hexyl	—	00B-4500-AN	00D-4500-AN	00F-4500-AN	AJO-8788

for 2.1 mm ID

1.7 µm MidBore Columns (mm)				SecurityGuard™ ULTRA Cartridges [‡]
Phases	30 x 3.0	50 x 3.0	100 x 3.0	3/pk
XB-C18	00A-4498-YO	00B-4498-YO	00D-4498-YO	AJO-8775
C18	—	00B-4475-YO	00D-4475-YO	AJO-8775
C8	00A-4499-YO	00B-4499-YO	00D-4499-YO	AJO-8777
HILIC	—	00B-4474-YO	—	AJO-8779

for 3.0 mm ID

1.3 µm Minibore Columns (mm)		
Phases	30 x 2.1	50 x 2.1
C18	00A-4515-AN	00B-4515-AN

[‡] SecurityGuard ULTRA Cartridges require holder, Part No.: AJO-9000

Terms and Conditions

Subject to Phenomenex Standard Terms & Conditions, which may be viewed at www.phenomenex.com/TermsAndConditions.

Trademarks

Phenomenex, Clarity, and Kinetex are registered trademarks of Phenomenex. Axia, Aeris, Oligo-MS, MidBore, Novum, Phenex, and SecurityGuard are trademarks of Phenomenex. Waters, ACQUITY, and XSelect are registered trademarks, and CSH is a trademark of Waters Corporation. Agilent and Poroshell are registered trademarks of Agilent Technologies, Inc. Ascentis and Supelco are registered trademarks of Sigma-Aldrich Co. LLC. ACE is a registered trademark and UltraCore is a trademark of Advanced Chromatography Technologies Limited. Accucore is a trademark and Hypersil GOLD is registered trademark of Thermo Hypersil-Keystone. API 4000 is a trademark of AB SCIEX Pte. Ltd.

Disclaimer

Phenomenex is not affiliated with Advanced Materials Technology, Inc., Agilent Technologies, Sigma-Aldrich Co., Thermo Hypersil-Keystone or Waters Corporation.

Comparative separations may not be representative of all applications.

SecurityGuard is patented by Phenomenex. U.S. Patent No. 6,162,362.

CAUTION: this patent only applies to the analytical-sized guard cartridge holder, and does not apply to SemiPrep, PREP or ULTRA holders, or to any cartridges.

Axia column and packing technology is patented by Phenomenex. U.S. Patent No. 7,674,383


Kinetex EVO is patented by Phenomenex. U.S. Patent Nos. 7,563,367 and 8,658,038 and foreign counterparts.

© 2015 Phenomenex, Inc. All rights reserved.

NEW Kinetex F5

HPLC/UHPLC Core-Shell Columns

- Reduce Method Development Time by Days
- Greater Reproducibility than other **PFPs**
- 5 Glorious Interaction Mechanisms
- 5 Valuable LC Separation Modes



Australia
t: +61 (0)2-9428-6444
f: +61 (0)2-9428-6445
auinfo@phenomenex.com

Austria
t: +43 (0)1-319-1301
f: +43 (0)1-319-1300
anfrage@phenomenex.com

Belgium
t: +32 (0)2 503 4015 (French)
t: +32 (0)2 511 8666 (Dutch)
f: +31 (0)30-2383749
beinfo@phenomenex.com

Canada
t: +1 (800) 543-3681
f: +1 (310) 328-7768
info@phenomenex.com

China
t: +86 (0)20 2282-6668
f: +86 (0)20 2809-8130
chinainfo@phenomenex.com

Denmark
t: +45 4824 8048
f: +45 4810 6265
nordicinfo@phenomenex.com

Finland
t: +358 (0)9 4789 0063
f: +45 4810 6265
nordicinfo@phenomenex.com

France
t: +33 (0)1 30 09 21 10
f: +33 (0)1 30 09 21 11
franceinfo@phenomenex.com

Germany
t: +49 (0)6021-58830-0
f: +49 (0)6021-58830-11
anfrage@phenomenex.com

India
t: +91 (0)40-3012 2400
f: +91 (0)40-3012 2411
indiainfo@phenomenex.com

Ireland
t: +353 (0)1 247 5405
f: +44 1625-501796
eireinfo@phenomenex.com

Italy
t: +39 051 6327511
f: +39 051 6327555
italiainfo@phenomenex.com

Luxembourg
t: +31 (0)30-2418700
f: +31 (0)30-2383749
nlinfo@phenomenex.com

Mexico
t: 001-800-844-5226
f: 001-310-328-7768
tecnicomx@phenomenex.com

The Netherlands
t: +31 (0)30-2418700
f: +31 (0)30-2383749
nlinfo@phenomenex.com

New Zealand
t: +64 (0)9-4780951
f: +64 (0)9-4780952
nzinfo@phenomenex.com

Norway
t: +47 810 02 005
f: +45 4810 6265
nordicinfo@phenomenex.com

Puerto Rico
t: +1 (800) 541-HPLC
f: +1 (310) 328-7768
info@phenomenex.com

Spain
t: +34 91-413-8613
f: +34 91-413-2290
espinfo@phenomenex.com

Sweden
t: +46 (0)8 611 6950
f: +45 4810 6265
nordicinfo@phenomenex.com

United Kingdom
t: +44 (0)1625-501367
f: +44 (0)1625-501796
ukinfo@phenomenex.com

USA
t: +1 (310) 212-0555
f: +1 (310) 328-7768
info@phenomenex.com

All other countries
Corporate Office USA 
t: +1 (310) 212-0555
f: +1 (310) 328-7768
info@phenomenex.com